

Brain abnormalities on neuroimaging in Children with Congenital Zika Syndrome in Salvador, Brazil, and its possible implications on neuropsychological development

Paula Sanders Pereira Pinto^{1,2}  | Thalita Madeira de Almeida²  |
Lucas Monteiro²  | Mirela Maisa da Silva Souza²  |
George Anderson Alves dos Santos²  | Cristiane Wanderley Cardoso³  |
Letícia Marques dos Santos⁴  | Guilherme Sousa Ribeiro^{5,6}  | Darci Neves dos Santos² 

¹UNIFACS, Universidade Salvador, Salvador, Brazil

²Instituto de Saúde Coletiva (UFBA), Salvador, Brazil

³Secretaria Municipal de Saúde de Salvador, Salvador, Brazil

⁴Instituto de Humanidades, Artes e Ciências da UFBA, Salvador, Brazil

⁵Fundação Oswaldo Cruz, Brazil

⁶Faculdade de Medicina da UFBA, Salvador, Brazil

Correspondence

Paula Pinto, UNIFACS, Universidade Salvador, AV. Luis Viana Filho, Salvador, Bahia 41770-235, Brazil.
Email: psanderspinto@gmail.com

Funding information

This research project was funded by CAPES and CNPq n° 440557/2016-0

Abstract

Objective: To characterize the spectrum of brain damages presented in children affected by Congenital Zika Syndrome (CZS), verify the existence of a co-occurrence pattern of these damages and discuss possible implications for the neuropsychological development.

Methods: Descriptive, quantitative, individualized, and cross-sectional study using secondary sources. We selected 136 children with CZS from the database of the Center of Strategic Information on Health Vigilance of the Municipal Office of Salvador, Brazil. We conducted descriptive and multiple correspondence analyses.

Results: Among the set of analyzed variables, microcephaly (51.5%), ventriculomegaly (57.4%), and brain calcifications (77.2%) were identified as the most frequent. The multiple correspondence analysis showed that the combination of these three variables (32.4%) was what better represented the spectrum of brain damages in the Central Nervous System.

Interpretation: Damage in the sensory-motor, cognitive and language development, as well as neurodevelopmental disorders, are described in the literature as impairments associated, either isolated or combined, with these damages, and it is worth highlighting that, in combined brain damages, impairments tend to be more severe. The findings of this study may contribute to understanding the repercussions of CZS on the neuropsychological development of children affected by the epidemic.

KEYWORDS

Congenital Zika Syndrome, neurological spectrum, neuropsychological development

1 | INTRODUCTION

Phylogenetic studies indicate that the Zika virus (ZIKV) was probably introduced in Brazil in 2013 (Campos, Bandeira, & Sardi, 2015; Faria et al., 2016), but only in

April 2015 it was first identified in the country, being recognized as the causative agent of a rash-disease outbreak in Brazil's Northeast region (Zanluca et al., 2015). Only in Salvador, the country's fourth largest city and one of the epicenters of the ZIKV epidemic in the Northeast, almost

15,000 cases of a rash disease attributed to ZIKV were reported between February and June 2015 (Cardoso et al., 2015).

Initially, ZIKV infections were considered of little relevance due to their self-limited and poorly symptomatic clinical picture. However, in the months following the peak of the ZIKV epidemic, there was an increase in records of children born with microcephaly, which was later associated (Paploski et al., 2016) with ZIKV infection during pregnancy (Awadh et al., 2017). Until the end of 2015, more than 3,000 babies with microcephaly, corresponding to 20 cases for every 10,000 births, were recorded in Brazil (Awadh et al., 2017). Later, it was observed that, besides microcephaly, children affected by an intrauterine ZIKV infection had a series of brain damages, as well as damages in other organs and systems, which led to the recognition of the Congenital Zika Syndrome (CZS) (Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Secretaria de Atenção à Saúde, 2017).

Congenital Zika Syndrome is characterized by serious and direct brain damages in brain volume, including structural and functional components observed through neuroimaging examinations of the Central Nervous System (CNS), such as transfontanelar ultrasound, computed tomography, and magnetic resonance (Moore et al., 2017). Are characteristics of the CZS: severe microcephaly with partially collapsed skull; thin cerebral cortices with subcortical calcifications; ocular changes, such as macular scarring and pigments in the center of the retina; arthrogryposis; as well as congenital contractures, early hypertonia and symptoms of extrapyramidal involvement, among others (Moore et al., 2017; Wheeler, 2018).

Although evidence suggests that ZIKV infections can occur during any gestation stage, associations with brain damages in the embryo or fetus are more frequent during the first trimester (Molnár & Kennedy, 2016) and, when the infection occurs during this time, there is an increased chance of abortion, stillbirth, early mortality, and microcephaly (Wheeler, 2018). Although it has been shown that the brain damages seen in this syndrome are associated with neurodevelopmental impairments, such as language and cognition deficits and problems in motor and behavioral skills (Alves, Paredes, Silva, Mello, & Alves, 2018; França et al., 2018), the literature on the subject is still scarce, hindering the understanding of the occurrence patterns of the brain damages, and the possible consequences for the neurodevelopment of children.

This study aimed to characterize the spectrum of brain damages detected by neuroimaging examination of children with confirmed CZS in Salvador, Bahia, Brazil, verify the existence of a co-occurrence pattern of these damages, and discuss the possible implications for the neuropsychological development of this generation.

2 | METHODS

2.1 | Study design

This is a descriptive, quantitative, individualized, and cross-sectional study.

2.2 | Sample selection

We selected the sample of this study from the database containing the case records of microcephaly and CZS provided by the Center of Strategic Information on Health Vigilance (Centro de Informações Estratégicas de Vigilância em Saúde, CIEVS) from Salvador's municipal office. This database was previously used to characterize CZS cases in Salvador and evaluate the accuracy of measuring the cephalic perimeter to detect children with CZS (Kikuti et al., 2018). During the use of the database in December 2016, there were 714 reports of children born in Salvador between August 01, 2015 and July 31, 2016. This period corresponds to that with the largest number of children born with suspected CZS (França et al., 2016). Among the reported cases, 433 (60.6%) had their investigation concluded by Center of Strategic Information on Health Vigilance, of which 47.1% (204/433) were confirmed. In November 2017, based on new investigations, 22 confirmed cases born in this period were included, totaling a sample of 224 children. Of these, 136 (60.7%) were selected with complete information on the relevant variables for this study. Of the remain, 38 (43.2%) were positive for STORCH (syphilis, toxoplasmosis, rubella, cytomegalovirus, herpes simplex) and 50 (56.8%) were missing data for variables of interest. In addition there was no difference between those who were selected or excluded. The mean for maternal age was 26.7 years (*SD* 6.35) for participants and 27.4 (*SD* 7.9) for the loss; gestational age correspond to 37 weeks (*SD* 3.2) and 36 (*SD* 4.1), respectively; and 58% of girls participating, against 63% not involved.

2.3 | Criteria for reporting CZS cases

The criteria used to report and confirm the cases included in this study followed protocols established by the Ministry of Health, which changed in the course of the CZS epidemic (Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância das Doenças Transmissíveis, 2016; Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância Epidemiológica, 2015; Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Secretaria de Atenção à Saúde, 2017).

2.4 | Criteria for excluding cases

We excluded, from this study, children that had positive laboratory results for any of the STORCH or had genetic alterations.

2.5 | Definition of the variables

The analyzed variables were: (a) Age of the mother; (b) Self-declared race; (c) Income of the mother; (d) Marital status of the mother; (e) Education level of the mother; (f) Number of children in the same gestation; (g) Prenatal care; (h) Number of prenatal visits; (i) Drinking and smoking habits during gestation; (j) Rash during gestation; (k) Pruritus during gestation; (l) Fever during gestation; (m) Trimester in which rash occurred; (n) Sex of the baby; (o) Type of delivery; (p) Weight at birth; (q) Apgar at 1 and 5 min after birth; (r) Time of detection of brain damages; (s) Occurrence of microcephaly according to the Intergrowth-21st table; (t) Calcifications; (u) Ventriculomegaly; (v) Subependymal cysts; (w) Dysgenesis of the corpus callosum; (x) Agenesis of the corpus callosum; (y) Hydrocephalus; (z) Lissencephaly.

2.6 | Analysis plan

To tabulate and organize the database, we used the software SPSS v. 25 IBM Corporation. For the descriptive analyses, we calculated absolute and relative frequencies. Aiming to identify co-occurrence patterns of some brain damages, we used a multiple correspondence analysis by means of the software R *Foundation for Statistical Computing*, the packages used were *FactoMineR* and *factoextra*. This analysis was conducted to explore the reduction of categorical variables, determining the degree of global association between observations and variables, and allowing to perceive possible associations between variables regarding some brain damages seen in the children's neuroimaging examinations (Infantosi, Costa, & Almeida, 2014; Nascimento, Almeida, Castilho, & Infantosi, 2013). The generated graphs are formed by crossing the two dimensions that better explain the data variability, each dimension consisting of the variables with the highest values of relative contribution. The graphs show the position of each response category so that these positions can be interpreted as associations between response patterns. Dots that appear together in the graph suggest a stronger association between the brain damages categories, forming groups that are displayed using oval diagrams (Infantosi et al., 2014).

2.7 | Ethical considerations

The research project "Effects of the congenital neurological manifestations associated with ZIKV on children

development: a prospective cohort study in the primary care context in Salvador-BA," which integrates this study, was approved by the Ethics Committee of the Institute of Collective Health of the Federal University of Bahia, under the number 1.659.107, meeting the requirements of the Resolution 466/12 of the National Health Council regarding research with human beings.

The children that took part in this study were inserted in a longitudinal prospective research study to follow child development with measurements of cognitive, language, and motor development evaluated through the Bayley-III Scale. They comprise the cohort DICa, in follow-up at the Institute of Collective Health of the Federal University of Bahia.

2.8 | Literature review

To discuss the probable implications of CZS for the neuropsychological development, we conducted an unsystematic literature review, through the snowball method using the Database of the CAPES Portal of articles published in the last 20 years. We used the following descriptors: "microcephaly and neurodevelopment"; "ventriculomegaly and neurodevelopment"; "calcifications and neurodevelopment."

3 | RESULTS

Among the 136 analyzed children, the median of the age of the mothers was 27 (15–44) years. Most mothers (116/136, 85.3%) were between 16 and 34 years old and 91.3% (115/126) declared themselves as black. Regarding family income, 60.6% (40/66) lived with less than one minimum wage per month. Regarding marital status, 54.2% were single, with 53.2% having concluded high school. Only 12.6% (14/111) had a college degree.

On gestation history, 94.7% (126/133) generated a single child, 98.4% (124/126) had prenatal care, with 71.1% (86/121) having more than six visits. A total of 12.1% (16/132) consumed alcohol or tobacco during the gestation. There were reports of symptoms suggesting ZIKV infection, such as rash (79/121, 65.3%), fever (45/115, 39.1%), and pruritus (61/121, 50.4%). Rash appeared mostly in the first gestational trimester (53/81, 65.4%).

Regarding the babies, 79 (79/136) were female, 28.6% (38/133) were premature, and 36.3% (49/135) had low weight at birth (<2500 g). Most had an adequate performance on Apgar (8–10), with 81.1% (107/132) 1 min and 96.2% (127/132) 5 min after birth.

Of the children confirmed for CZS, 64% had brain damages detected in the postpartum period (87/101, 64%). The cephalic perimeter of 48.5% of the children was within the normal range, 19.1% were diagnosed with microcephaly, and

32.4% with severe microcephaly. Other brain damages are described in Table 1.

Sixty-eight children had between three and five of the brain damages in the CNS included in this study. The others had fewer than three damages. The multiple correspondence analysis was initially performed with six variables regarding the brain damages that were present in more than 10% of the 136 children (ventriculomegaly, hydrocephalus, calcifications, subependymal cysts, microcephaly, and dysgenesis of the corpus callosum). Based on the results, the variables hydrocephalus and subependymal cysts were excluded after the first level of analysis, indicating a relative contribution to inertia (variability) well below the usually recommended limit ($1/Q$, in which Q is the number of variables) (Nascimento, 2013). The examination of the remaining variables allowed to identify the contribution of ventriculomegaly, calcifications, and microcephaly for dimension 1, suggesting a co-occurrence pattern (Table 2). Dimension 2, consisting only of dysgenesis, explained only 25.3% of the data variability and was used only for comparison, while dimension 1 explained 39.9% (Table 2 and Figure 1). The Figure 1 shows the main brain damages in the children of this study. Based on the results provided by the correspondence analysis, 44 (44/136, 32.4%) participants had co-occurrence of the three variables. The Figure 2 indicate a significant difference between the categories of all variables.

TABLE 1 Characteristics of the brain damages of children with confirmed CZS, born between 01 August 2015 and 31 July 2016 in Salvador, BA, Brazil

Characteristics	<i>n</i>	%
Period of detection of brain damage (135)		
Intrauterine	48	35.6
Postpartum	87	64.4
Classification according to cephalic perimeter (136)		
Within normal range	66	48.5
Microcephaly ($\leq -2SD$ to $< -3SD$)	26	19.1
Severe microcephaly ($\leq -3SD$)	44	32.4
Brain damages (136)		
Calcifications	105	77.2
Ventriculomegaly	78	57.4
Subependymal cysts	19	13.9
Dysgenesis of the corpus callosum	16	11.8
Agenesis of the corpus callosum	14	10.3
Hydrocephalus	15	11.0
Lissencephaly	10	7.4

Source: Registro de Eventos em Saúde Pública/ Secretaria Municipal de Saúde de Salvador/Centro de Informações Estratégicas em Vigilância em Saúde.

4 | DISCUSSION

The brain damages identified among the participants of this study confirm previous studies regarding the similarity of the damages in children affected by CZS (Moore et al., 2017; Wheeler, 2018). The multiple correspondence analysis identified a global association between microcephaly, ventriculomegaly, and calcifications, indicating a probable pattern of co-occurrence of changes in CZS. Combined brain damages are frequent in congenital syndromes and increase the degree of neurodevelopmental impairments (Gaglioti et al., 2005; Hetts, Sherr, Chao, Gobuty, & Barkovich, 2006). While more frequent impairments include cerebral palsy, epilepsy, intellectual disability, language impairments, difficulty swallowing, and anomalies of the visual and auditory systems. Attention-deficit/hyperactivity disorder (ADHD) and autistic spectrum disorder (ASD) were highlighted among the neurodevelopmental disorders (Brunoni et al., 2016; Wheeler, 2018), but there were babies who developed normally despite the presence of these abnormalities (Arroyo, 2018; WHO, 2016). Microcephaly was absent in almost half of the studied children. However, other brain damages are also frequent in CZS (França et al., 2016; Moore et al., 2017), even in the absence of microcephaly (Van der Linden et al., 2016).

Microcephaly has a role in determining the spectrum of neuropsychological and brain damages, as it is known that the smaller the cephalic perimeter the more serious the clinical picture presented by the child. Regarding our finding of 51.5% in cephalic perimeter change, it is likely that these children have a more prominent neurological impairment when compared to those without microcephaly. A risk of 10.5% for the occurrence of intellectual disability is estimated when the cephalic perimeter is between 2 and 3 standard deviations below the expected mean, increasing to 51.2% with a perimeter between 3 and 4 standard deviations and reaching 100% for those with 4 standard deviations below the mean (Alves et al., 2018). A study with 82 children with cerebral palsy, probably related to CZS, used the Bayley-III scale to evaluate cognitive performance and found an association between very low cognitive performance and birth and follow-up cephalic perimeter, as well as between the motor score and follow-up cephalic perimeter (Carvalho et al., 2019). The neuropsychological impairments differ according to cause, affected brain area and extension. A cephalic perimeter smaller than 2 standard deviations below the mean can be found in about 2% of the population but most have a normal intellectual performance (Arroyo, 2018).

Regarding CVS, studies revealed the possibility of microcephaly occurring after the first months of life in children that were born with an adequate cephalic perimeter, indicating that the virus' replication may continue in newborns (Van der Linden et al., 2016). Cognitive

TABLE 2 Description of the remaining brain damages variables and percentage of relative contribution values for variability in dimensions 1 and 2 obtained through multiple correspondence analysis

Variable	Categories [%]	Variability of relative contribution (%)	
		Dimension 1	Dimension 2
Ventriculomegaly	Sim [57.4]	27.32	18.14
Calcifications	Sim [77.2]	30.25	5.82
Dysgenesis	Sim [11.8]	8.71	68.91
Microcephaly	Sim [51.5]	33.73	7.13

FIGURE 1 Correspondence map for a multiple correspondence analysis activities variables. Correspondence map with the categories of Calcifications, Dysgenesis, Microcephaly, and Ventriculomegaly. Variable categories with a similar profile are grouped together (absence = 0 and presence = 1) [Colour figure can be viewed at wileyonlinelibrary.com]




FIGURE 2 Confidence ellipses round the categories of activities variables. Confidence ellipses (95%) for categories of Calcifications, Dysgenesis, Microcephaly, and Ventriculomegaly. The objective is to see whether the categories of a categorical variable are significantly different from each other. Categories without ellipses intersection are significantly different (absence = 0 and presence = 1) [Colour figure can be viewed at wileyonlinelibrary.com]

impairments in the first year may be absent among individuals that acquired microcephaly after birth or showed no other congenital anomaly at birth; however, impairments may appear during childhood or school years, requiring

special educational and emotional support (Subissi et al., 2018; Wheeler, 2018).

Sensory disabilities, when occurring together with microcephaly, may make the development even more difficult,

because they reduce the capacity to learn and develop with the environment to which one belongs. Studies have indicated that 1-year-old babies may show a cognitive development equivalent to that of 2- or 3-month-old babies, the worst performance being that of language (Alves et al., 2018).

Ventriculomegaly occurred in 57.4% of the investigated children. This is one of the most frequent impairments in congenital syndromes, characterized by the dilation of the cerebral ventricles and, although isolated from other brain damages, can generate delays, especially on motor and cognitive development (Scala et al., 2017). A meta-analysis was conducted with 108 children that presented isolated unilateral ventriculomegaly from congenital or genetic changes and seven were diagnosed with neurodevelopmental delay. The neuropsychological impairments were associated with different degrees of ventriculomegaly (mild: 10 to 12 mm; moderate: 12.1 to 14.9 mm; severe: ≥ 15 mm), as well as with morphological or chromosomal anomalies and with the male sex (Scala et al., 2017).

It has been shown that the milder the degree of ventriculomegaly, the higher the probability of reaching a typical neurodevelopment (Gaglioti et al., 2005; Goldstein, Copel, & Makhoul, 2005), especially with a normal karyotype (Goldstein et al., 2005) and if the ventriculomegaly is isolated from other brain damages (Goldstein et al., 2005; Nascimento, 2013). The atrial size considered the threshold (between 10 and 11 mm) (Gaglioti et al., 2005) was also considered a good prognosis for the development of children with ventriculomegaly. Brain calcifications together with ventricular anomalies and microcephaly are common in congenital infections. Our study found 77.2% of brain calcifications. An investigation with 22 children whose mothers had toxoplasmosis during the pregnancy identified 72.7% of isolated calcifications or combined with other brain damages, being 40.9% with isolated calcifications, 9% combined with ventricular dilations, 4.5% with mild increase of the ventricles and the subarachnoid cisterns, 4.5% with porencephalic cyst, 4.5% with hydrocephalus, and 4.5% combined with hydranencephaly (Melamed, Dornelles, & Eckert, 2001). In CZS, neuroimaging examinations have detected predominantly calcifications of diffuse and punctiform nature, mainly in the cortical-subcortical junction of the frontal and parietal lobes (Carvalho et al., 2019), as well as areas such as the brainstem, basal nuclei, periventricular region and thalamus (Martins, 2017). In a study conducted with four infants up to 4 months old with presumable congenital ZIKV infection, having calcifications in the cortical-subcortical junction and ventriculomegaly, all participants were observed to have hyperreflexia and hypertonia, atypical development and deficit in hand function. The authors discuss the correlation of the outcomes with the referred brain damages (Botelho, Neri, & Silva, 2016).

From the results of our study and the scientific evidence, it is supposed that the isolated brain damages or, especially,

those that exist in a co-occurrence pattern consisting of ventriculomegaly, calcifications, and microcephaly, are associated with a set of neurodevelopmental impairments in children with CZS.

The level of impairment of these injuries has already been indicated by some studies (Alves et al., 2018; Carvalho et al., 2019; França et al., 2018). After following 24 children with an average age of 19.9 months and confirmed diagnosis for CZS in Pernambuco, Brazil, a study (Alves et al., 2018) verified that they presented a developmental level equivalent to 2.1-month-old children for language, 2.7 month old for global motricity, 3.1 month old for fine motricity and 3.4 month old for psychosocial features, evaluated by the Denver-II test. The children did not walk or speak, which is expected from their age. Another study (França et al., 2018) evaluated eight children with CSZ in Rio Grande do Norte, Brazil through the Bayley-III scale and compared them with typical children, both with an average age of 20.5 months. The authors found a significant delay in cognitive development, as well as in the motor domain in comparison with typical children. A third study (Carvalho et al., 2019) found a very low performance in the cognitive, motor, and language performances of children with cerebral palsy probably associated with CZS.

One of the limitations of this study is the loss of 40% of subjects in the case selection due to incomplete data for the variables of interest, as well as those confirmed for STORCH, because these infections were not the object of this study. However, no relevant differences for child sex, maternal age, and gestational age at birth were observed between those who were selected or excluded.

In addition, the case definition for CZS supported by neuroimaging has not been confirmed by ZIKV serology, weakening the association between neurological findings and maternal ZIKV infection. However, there was no availability of adequate tests to detect the virus at the time. Although the use of secondary data offers limitations, to our knowledge this research is the first to discuss data from the epidemiological surveillance center during the outbreak on the perspective of child development.

5 | CONCLUSIONS

This study contributes to the knowledge of CZS and its spectrum of most frequent brain damages, identifying a co-occurrence pattern among these damages and discussing the possible effects on the neuropsychological development of children. The existence of a combination of brain damages tends to promote more severe impairments than those found when they occur in isolation. Thus, in any dimension of child development, adequate care needs to be offered, especially in an early stage.

ACKNOWLEDGEMENTS

We thank the CAPES and CNPQ funders and the Ministry of Health for support, as well as the Professor Dr. Rosemeire Fiaconne for the statistical support in the data analysis.

CONFLICT OF INTEREST

We declare that no author has a conflict of interest.

ETHICS APPROVAL STATEMENT

The research project, which integrates this study, was approved by the Ethics Committee of the Institute of Collective Health of the Federal University of Bahia, under the number 1.659.107.

AUTHOR CONTRIBUTIONS

Paula Sanders Pereira Pinto and Darci Neves dos Santos: planning, substantial contributions for the research strategy, acquisition, analysis and interpretation of the data; writing and review of the manuscript; approval of the submitted and final versions; Thalita Madeira de Almeida; Lucas Monteiro; Mirela Maisa da Silva Souza; George Anderson Alves dos Santos; Cristiane Wanderley Cardoso; Letícia Marques dos Santos: substantial contributions for the research strategy, acquisition, analysis and interpretation of the data; writing and review of the manuscript; approval of the submitted and final versions; Guilherme Sousa Ribeiro: writing and critical review of the article; approval of the submitted and final versions.

DATA AVAILABILITY STATEMENT

Author elects to not share data.

ORCID

Paula Sanders Pereira Pinto  <https://orcid.org/0000-0002-8824-4688>

Thalita Madeira de Almeida  <https://orcid.org/0000-0002-9257-1278>

Lucas Monteiro  <https://orcid.org/0000-0002-4832-7059>

Mirela Maisa da Silva Souza  <https://orcid.org/0000-0001-6056-050X>

George Anderson Alves dos Santos  <https://orcid.org/0000-0002-5600-5287>

Cristiane Wanderley Cardoso  <https://orcid.org/0000-0001-5432-1174>

Letícia Marques dos Santos  <https://orcid.org/0000-0001-5963-2166>

Guilherme Sousa Ribeiro  <https://orcid.org/0000-0002-6798-2059>

Darci Neves dos Santos  <https://orcid.org/0000-0003-1500-8294>

REFERENCES

Alves, L. V., Paredes, C. E., Silva, G. C., Mello, J. G., & Alves, J. G. Neurodevelopment of 24 children born in Brazil with Congenital

Zika Syndrome in 2015: A case series study. *British Medical Journal Open*, 2018, 8(e021304), 1–5. <https://doi.org/10.1136/bmjopen-2017-021304>

- Arroyo, H. A. (2018). Microcefalia. *Medicina*, 78(Suppl. II), 94–100.
- Awadh, A., Chughtai, A. A., Dyda, A., Sheikh, M., Heslop, D. J., & MacIntyre, C. R. (2017). Does Zika virus cause microcephaly—Applying the Bradford Hill viewpoints. *PLoS Currents Outbreaks*, 22(1), 1–11. <https://doi.org/10.1371/currents.outbreaks.2fced6e886074f6db162a00d4940133b>
- Botelho, A. C. G., Neri, L. V., & Silva, M. Q. F. (2016). Infecção congênita presumível por Zika vírus: Achados do desenvolvimento neuropsicomotor—Relato de casos. *Revista Brasileira De Saúde Materno Infantil*, 16(Supl. 1), 45–50. <https://doi.org/10.1590/1806-9304201600S100004>
- Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância das Doenças Transmissíveis. (2016). *Protocolo de vigilância e resposta à ocorrência de microcefalia e/ou alterações do sistema nervoso central (SNC)* / Ministério da Saúde, Secretaria de Vigilância em Saúde, Departamento de Vigilância das Doenças Transmissíveis. Brasília: Ministério da Saúde.
- Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância Epidemiológica. (2015). *NOTA INFORMATIVA Nº 01/2015—COES MICROCEFALIAS* / Ministério da Saúde. Secretaria de Vigilância em Saúde. Departamento de Vigilância Epidemiológica. Brasília: Ministério da Saúde.
- Brasil. Ministério da Saúde. Secretaria de Vigilância em Saúde. Secretaria de Atenção à Saúde. (2017). *Orientações integradas de vigilância e atenção à saúde no âmbito da Emergência de Saúde Pública de Importância Nacional: Procedimentos para o monitoramento das alterações no crescimento e desenvolvimento a partir da gestação até a primeira infância, relacionadas à infecção pelo vírus Zika e outras etiologias infecciosas dentro da capacidade operacional do SUS [electronic resource]* / Ministério da Saúde, Secretaria de Vigilância em Saúde, Secretaria de Atenção à Saúde. Brasília: Ministério da Saúde.
- Brunoni, D., Blascovi-Assis, S. M., Osório, A. A. C., Seabra, A. G., Amato, C. A. D. L. H., Teixeira, M. C. T. V., ... Carreiro, L. R. R. (2016). Microcefalia e outras manifestações relacionadas ao vírus Zika: impacto nas crianças, nas famílias e nas equipes de saúde. *Ciência & Saúde Coletiva*, 21(10), 3297–3302. <https://doi.org/10.1590/1413-812320152110.16832016>
- Campos, G. S., Bandeira, A., & Sardi, S. (2015). Zika virus outbreak, Bahia Brazil. *Emerging Infectious Diseases*, 21(10), 1885–1886. <https://doi.org/10.3201/eid2110.150847>
- Cardoso, C. W., Paploski, I. A. D., Kikuti, M., Rodrigues, M. S., Silva, M. M. O., Campos, G. S., ... Ribeiro, G. S. (2015). Outbreak of exanthematous illness associated with Zika, Chikungunya, and Dengue viruses, Salvador, Brazil. *Emerging Infectious Diseases*, 21(12), 2274–2276. <https://doi.org/10.3201/eid2112.151167>
- Carvalho, A., Brites, C., Mochida, G., Ventura, P., Fernandes, A., Lage, M. L., ... Lucena, R. (2019). Clinical and neurodevelopmental features in children with cerebral palsy and probable Congenital Zika. *Brain and Development*, 41(7), 587–594. <https://doi.org/10.1016/j.braindev.2019.03.005>
- Faria, N. R., Azevedo, R. D. S. D. S., Kraemer, M. U. G., Souza, R., Cunha, M. S., Hill, S. C., ... Vasconcelos, P. F. C. (2016). Zika virus in the Americas: Early epidemiological and genetic findings. *Science*, 352(6283), 345–349. <https://doi.org/10.1126/science.aaf5036>
- França, G. V. A., Schuler-Faccini, L., Oliveira, W. K., Henriques, C. M. P., Carmo, E. H., Pedi, V. D., ... Victora, C. G. (2016). Congenital

- Zika virus syndrome in Brazil: A case series of the first 1501 live-births with complete investigation. *The Lancet*, 388, 891–897. [https://doi.org/10.1016/S0140-6736\(16\)30902-3](https://doi.org/10.1016/S0140-6736(16)30902-3)
- França, T., Medeiros, W., Souza, N., Longo, E., Pereira, S., França, T., & Sousa, K. (2018). Growth and development of children with microcephaly associated with Congenital Zika virus syndrome in Brazil. *International Journal of Environmental Research and Public Health*, 15(9), 1–11. <https://doi.org/10.3390/ijerph15091990>
- Gaglioti, P., Danelon, D., Bontempo, S., Mombro, M., Cardaropoli, S., & Todros, T. (2005). Fetal cerebral ventriculomegaly: Outcome in 176 cases. *Ultrasound in Obstetrics and Gynecology*, 25(4), 372–377. <https://doi.org/10.1002/uog.1857>
- Goldstein, I., Copel, J. A., & Makhoul, I. R. (2005). Mild cerebral ventriculomegaly in fetus: Characteristics and outcome. *Fetal Diagnosis and Therapy*, 20(4), 281–284. <https://doi.org/10.1159/000085086>
- Hetts, S. W., Sherr, E. H., Chao, S., Gobuty, S., & Barkovich, A. J. (2006). Anomalies of the corpus callosum: An MR analysis of the phenotypic spectrum of associated malformations. *American Journal of Roentgenology*, 187(5), 1343–1348. <https://doi.org/10.2214/AJR.05.0146>
- Infantosi, A. F. C., Costa, J. C. G. D., & Almeida, R. M. V. R. (2014). Análise de Correspondência: Bases teóricas na interpretação de dados categóricos em Ciências da Saúde. *Caderno De Saúde Pública*, 30(3), 473–486. <https://doi.org/10.1590/0102-311X00128513>
- Kikuti, M., Cardoso, C. W., Prates, A. P. B., Paploski, I. A. D., Kitron, U., Reis, M. G., ... Ribeiro, G. S. (2018). Congenital brain abnormalities during a Zika virus epidemic in Salvador, Brazil, April 2015 to July 2016. *Eurosurveillance*, 23(45), <https://doi.org/10.2807/1560-7917.ES.2018.23.45.1700757>
- Martins, M. R. (2017). *Teratogenicidade a Zika—um novo síndrome congênito? Faculdade de medicina de Lisboa: Trabalho final do Mestrado integrado em medicina*. Universidade de Lisboa. Retrieved from <http://repositorio.ul.pt/bitstream/10451/32609/1/MarianaRMartins.pdf>
- Melamed, J., Dornelles, F., & Eckert, G. (2001). Alterações tomográficas cerebrais em crianças com lesões oculares por toxoplasmose congênita. *Jornal De Pediatria*, 77(6), 475–480. <https://doi.org/10.1590/S0021-75572001000600010>
- Molnár, Z., & Kennedy, S. (2016). Neurodevelopmental disorders: Risks of Zika virus during the first trimester of pregnancy. *Nature Reviews Neurology*, 12, 315–316. <https://doi.org/10.1038/nrneuro.2016.71>
- Moore, C. A., Staples, J. E., Dobyns, W. B., Pessoa, A., Ventura, C. V., Fonseca, E. B. D., ... Rasmussen, S. A. (2017). Characterizing the pattern of anomalies in Congenital Zika Syndrome for pediatric clinicians. *JAMA Pediatrics*, 171(3), 288–295. <https://doi.org/10.1001/jamapediatrics.2016.3982>
- Nascimento, A. D. O., Almeida, R. M. V. R., Castilho, S. R., & Infantosi, A. F. C. (2013). Análise de correspondência múltipla na avaliação de serviços de farmácia hospitalar no Brasil. *Caderno de Saúde Pública*, 29(6), 1161–1172.
- Paploski, I. A. D., Prates, A. P. B., Cardoso, C. W., Kikuti, M., Silva, M. M. O., Waller, L. A., ... Ribeiro, G. S. (2016). Time lags between exanthematous illness attributed to Zika virus, Guillain-Barré Syndrome, and Microcephaly, Salvador, Brazil. *Emerging Infectious Diseases*, 22(8), 1438–1444. <https://doi.org/10.3201/eid2208.160496>
- Scala, C., Familiari, A., Pinas, A., Papageorghiou, A. T., Bhide, A., Thilaganathan, B., & Khalil, A. (2017). Perinatal and long-term outcomes in fetuses diagnosed with isolated unilateral ventriculomegaly: Systematic review and meta-analysis. *Ultrasound in Obstetrics and Gynecology*, 49, 450–459. <https://doi.org/10.1002/uog.15943>
- Subissi, L., Dub, T., Besnard, M., Maritéragi-Helle, T., Nhan, T., Lutringer-Magnin, D., ... Mallet, H.-P. (2018). Zika Virus infection during pregnancy and effects on early childhood development, French Polynesia, 2013–2016. *Emerging Infectious Diseases*, 24(10), 1850–1858. <https://doi.org/10.3201/eid2410.172079>
- van der Linden, V., Pessoa, A., Dobyns, W., Barkovich, A. J., Júnior, H. V. D. L., Filho, E. L. R., ... Moore, C. A. (2016). Description of 13 infants born during October 2015–January 2016 with Congenital Zika virus infection without microcephaly at birth—Brazil. *Morbidity and Mortality Weekly Report*, 65(47), 1343–1348. <https://doi.org/10.15585/mmwr.mm6547e2>
- Wheeler, A. C. (2018). Development of infants with Congenital Zika syndrome: What do we know and what can we expect? *Pediatrics*, 141(s2), 154–160. <https://doi.org/10.1542/peds.2017-2038D>
- WHO. (2016). *Screening, assessment and management of neonates and infants with complications associated with Zika virus exposure in utero*. Rapid Advice Guideline. Retrieved from <http://www.who.int/csr/resources/publications/zika/assessment-infants/en/>
- Zanluca, C., Melo, V. C., Mosimann, A. L. P., Santos, G. I. V., Santos, C. N. D., & Luz, K. (2015). First report of autochthonous transmission of Zika virus in Brazil. *Memórias do Instituto Oswaldo Cruz*, 110, 569–572.

How to cite this article: Sanders Pereira Pinto P, de Almeida TM, Monteiro L, et al. Brain abnormalities on neuroimaging in Children with Congenital Zika Syndrome in Salvador, Brazil, and its possible implications on neuropsychological development. *Int J Dev Neurosci*. 2020;80:189–196. <https://doi.org/10.1002/jdn.10016>